

Solution of in-class exercise 1: Bounding a sequence

Using the usual approach of subtracting the recurrence for $n - 1$, we obtain that for $n \geq 3$,

$$x_n - x_{n-1} = \sqrt{n} - \sqrt{n-1} + 2x_{n-1}$$

holds and therefore

$$x_n = (\sqrt{n} - \sqrt{n-1}) + 3x_{n-1}.$$

Now since $x_1 > 0$ we deduce that $x_n > 0$ for all n , and then in particular

$$x_n > 3^{n-1}x_1 \notin \mathcal{O}(2^n).$$

So the claim is **FALSE**.

Solution of in-class exercise 2: Random permutations

We prove the lemma by induction on the size n of the insertion permutation (or equivalently, the resulting tree).

Induction base case. If $n = 0$ or $n = 1$, the lemma trivially holds.

Induction step. Let $n \geq 2$ and suppose that the lemma holds for all insertion permutations of size strictly less than n . Let $\pi = (\pi(1), \dots, \pi(n))$ be a permutation drawn uniformly at random from \mathfrak{S}_n . The first element $\pi(1)$ will become the root of the tree T_π . Since the distribution of $\pi(1)$ is u.a.r. from $[n]$, the root of the tree is chosen uniformly at random, as required by the construction of $\tilde{\mathcal{B}}_{[n]}$.

Now let us tackle the two subtrees of the root: Let $k \in [n]$. We want to show that, conditioned on $\pi(1) = k$, the distribution of the left subtree of the root is the same as $\tilde{\mathcal{B}}_{\{1, \dots, k-1\}}$, and the distribution of the right subtree of the root is the same as $\tilde{\mathcal{B}}_{\{k+1, \dots, n\}}$. To this end, let π^- be the sequence of elements in π smaller than k , and let π^+ be the sequence of elements in π larger than k . Note that the insertion sequence will send the keys in π^- (π^+) in exactly this order to the left (right) subtree of the root. Since π^- and π^+ are uniformly random permutations of their respective element sets $\{1, \dots, k-1\}$ and $\{k+1, \dots, n\}$, we can apply the induction hypothesis to obtain that the left and right subtrees of the root are distributed just as we stated.

We conclude that this process produces a tree T_π that is distributed like $\tilde{\mathcal{B}}_{[n]}$.

Solution 1: A Random Tree? How Random?

According to Lemma 1.1 in the script, the probability of the tree is

$$\frac{1}{2 \cdot 1 \cdot 7 \cdot 1 \cdot 3 \cdot 1 \cdot 4} = \frac{1}{168}.$$

What do the extremal examples for the probability of a tree look like? In the case of 7 nodes, we consider the trees of height $7 - 1 = 6$ (which yields the smallest probability $\frac{1}{7!} = \frac{1}{5040}$) and the perfectly balanced tree of height 2 (which yields the largest probability $\frac{1}{7 \cdot 3^2 \cdot 1^4} = \frac{1}{63}$).

Since the problem on 7 nodes is finite, we could list all cases as a proof. However, there are 429 search trees on 7 nodes which would make the proof lengthy.

Alternatively, for the case of smallest probability, we can argue using Exercise 1.5 from the script (which is the same as in-class exercise 2):

Lemma 1. *If keys are inserted in a uniformly random order, the resulting search tree is distributed like \tilde{B}_s .*

It follows from the lemma that every random search tree is generated by at least one permutation on n elements (the probability of a random search tree is actually the number of permutations generating it divided by $n!$). Thus, the probability of a random search tree is at least $1/n!$ and since the extremal example of height $n - 1$ achieves exactly this probability, it has smallest probability. This holds for all n .

For the case of largest probability, let p_n denote the maximum probability for a random search tree with n vertices. Then we get the following recursion beginning with, by convention, $p_0 = 1$ (where k denotes the rank of the root).

$$p_n = \frac{1}{n} \cdot \max_{1 \leq k \leq n} (p_{k-1} p_{n-k}).$$

The first few values for this maximum probability can easily be computed, namely $p_1 = 1, p_2 = \frac{1}{2}, p_3 = \frac{1}{3}, p_4 = \frac{1}{8}, p_5 = \frac{1}{15}, p_6 = \frac{1}{36}, p_7 = \frac{1}{63}$. This already proves that the perfectly balanced tree on 7 nodes has indeed largest probability. To generalize this case to larger n , one would need to work somewhat harder.

Solution 2: Very Deep Nodes

Let N_{deep} denote the number of nodes of depth $n - 1$. We observe that a binary search tree B for n vertices has one node of depth $n - 1$ if and only if it is a path of length $n - 1$. Let p_n denote the probability that there is a node of depth $n - 1$. We have

$$\mathbb{E}[N_{\text{deep}}] = p_n \cdot 1 + (1 - p_n) \cdot 0 = p_n.$$

So it remains to compute p_n . Clearly, $p_1 = 1$. For $n \geq 2$ we apply induction. Note that if a tree T is a path then its root is either the smallest or the largest key. Hence

$$\begin{aligned} p_n &= \underbrace{\Pr[\text{rk}(\text{root}) = 1]}_{\frac{1}{n}} \cdot \underbrace{\Pr[\text{one node has depth } n-1 | \text{rk}(\text{root}) = 1]}_{p_{n-1}} + \\ &\quad \underbrace{\Pr[\text{rk}(\text{root}) = n]}_{\frac{1}{n}} \cdot \underbrace{\Pr[\text{one node has depth } n-1 | \text{rk}(\text{root}) = n]}_{p_{n-1}} \\ &= \frac{2}{n} \cdot p_{n-1} \end{aligned}$$

Induction yields that $p_n = \frac{2^{n-1}}{n!} \cdot p_1 = \frac{2^{n-1}}{n!}$ and so we are done.

REMARK. Note that this also provides us with the number of trees on n nodes that have height $n-1$: since each such tree is a path of length $n-1$, for each such tree there is exactly one ordering of n keys which produces this search tree (since a parent must always be inserted before its child). So each tree of height $n-1$ separately has a probability of $1/n!$. Since in total $p_n = 2^{n-1}/n!$, we conclude that there must be exactly 2^{n-1} trees of this type.

Of course, we could have found this number more easily: the number of trees on n nodes that are a single path is 2^{n-1} simply because for each edge (of which there are $n-1$ many) we can decide whether it should point to the left or to the right.

Solution 3: High Trees

By an (n, d) -tree we denote a tree on n vertices of height d . Let M_n denote the number of $(n, n-2)$ -trees and let M'_n denote the number of $(n, n-1)$ -trees. Our goal is to compute M_n . Clearly, $M_1 = 0$, $M_2 = 0$ and $M_3 = 1$. Now for $n \geq 4$, let us consider two ways to proceed.

Variante 1: by induction. Note that if a tree T in $\mathcal{B}_{\{1, \dots, n\}}$ has height $n-2$ then

- *either* the root of T is in $\{1, n\}$ and the subtree of the root is an $(n-1, n-3)$ -tree
- *or* the root of T is in $\{2, n-1\}$ and one of the subtrees of the root is an $(n-2, n-3)$ -tree.

As $n \geq 4$, the numbers $1, 2, n-1, n$ are distinct. So for $n \geq 4$,

$$M_n = 2 \cdot M_{n-1} + 2 \cdot M'_{n-2}, \tag{1}$$

As a direct consequence of Exercise 3 [Remark] we have $M'_n = 2^{n-1}$.

Equation (1) then yields that for $n \geq 4$,

$$M_n = 2 \cdot M_{n-1} + 2 \cdot 2^{n-3} = 2 \cdot M_{n-1} + 2^{n-2}.$$

By induction we obtain that $M_n = 2^i \cdot M_{n-i} + i \cdot 2^{n-2}$ for $i \leq n-3$. Hence for $n \geq 4$.

$$M_n = 2^{n-3} \cdot M_3 + (n-3) \cdot 2^{n-2} = 2^{n-3} + (n-3) \cdot 2^{n-2} = (2n-5) \cdot 2^{n-3}.$$

Variante 2: direct counting. We can count the number of trees on n vertices of height $n-2$ directly for $n \geq 3$: if a tree T in $\mathcal{B}_{\{1\dots n\}}$ has height $n-2$ then it consists of a path of height $n-2$ plus a leaf that can be attached at any of the first $n-2$ (out of the $n-1$) vertices of that path.

From Exercise 2 (1.9), we know that the number of paths on $n-1$ vertices is 2^{n-2} for $n \geq 1$. Multiplying by the number of possibilities for attaching the additional leaf, we obtain $(n-2)2^{n-2}$.




Figure 1: Trees that are counted twice.

This number, however, counts a certain kind of trees twice: All trees that have two leaves of depth $n-2$ (cf. Figure 1). How many such trees are there? Such a tree consists of a path on $n-2$ vertices plus two leaves attached to its end, so there are equally many of them as there are paths on $n-2$ vertices. Again using Exercise 2 (1.9), we obtain 2^{n-3} for this number. So in total, we obtain $(n-2)2^{n-2} - 2^{n-3}$ for $n \geq 3$ (which is the same as we obtained in Variante 1).

Solution 4: Solving Recurrences

- (1) First we compute that $a_1 = 1$ and $a_2 = \frac{3}{2}$. Now for $n \geq 3$, we multiply the recurrence relation by n and write it once for n and once for $n-1$. This yields

$$na_n = n + \sum_{i=1}^{n-1} a_i \tag{2}$$

and

$$(n-1)a_{n-1} = (n-1) + \sum_{i=1}^{n-2} a_i. \tag{3}$$

Now subtracting (3) from (2), we obtain

$$na_n - (n-1)a_{n-1} = 1 + a_{n-1}$$

and thus

$$a_n = \frac{1}{n} + a_{n-1}.$$

This recursion can easily be telescoped from which we obtain

$$a_n = \frac{1}{n} + \frac{1}{n-1} + \dots + \frac{1}{3} + \underbrace{a_2}_{=1/2+1} = H_n.$$

Therefore, $a_n = H_n$ for all $n \in \mathbf{N}$.

(2) We first compute that $b_1 = 1$ and $b_2 = 3$. Now for $n \geq 3$

$$b_n = 2 + \sum_{i=1}^{n-1} b_i \quad (4)$$

and

$$b_{n-1} = 2 + \sum_{i=1}^{n-2} b_i \quad (5)$$

Now subtracting (5) from (4), we obtain

$$b_n - b_{n-1} = b_{n-1},$$

therefore

$$b_n = 2 b_{n-1}$$

and thus

$$b_n = 2^{n-2} b_2 = 3 \cdot 2^{n-2}.$$

Therefore, $b_1 = 1$ and $b_n = 3 \cdot 2^{n-2}$ for all $n \geq 2$.

(3) We first compute that $c_0 = 0$ and $c_1 = 0$. Then for $n \geq 2$, we first note that

$$\sum_{i=1}^n \frac{c_{i-1} + c_{n-i}}{2} = \sum_{i=1}^n c_{i-1}$$

which then allows us to write the simpler recurrences for c_n and c_{n-1}

$$c_n = n - 1 + \sum_{i=1}^n c_{i-1} \quad (6)$$

and

$$c_{n-1} = n - 2 + \sum_{i=1}^{n-1} c_{i-1} \quad (7)$$

If we now subtract (7) from (6), then

$$c_n - c_{n-1} = 1 + c_{n-1}$$

and thus

$$c_n = 1 + 2 c_{n-1}.$$

For telescoping, it turns out to be convenient to divide the recurrence by 2^n , then we have

$$\frac{c_n}{2^n} = \frac{1}{2^n} + \frac{c_{n-1}}{2^{n-1}}$$

and we can telescope for $c_n/2^n$, yielding

$$\frac{c_n}{2^n} = \frac{1}{2^n} + \frac{1}{2^{n-1}} + \cdots + \frac{1}{2^2} + \frac{c_1}{2^1} = \frac{1}{2} - \frac{1}{2^n}.$$

Therefore, $c_0 = 0$ and $c_n = 2^{n-1} - 1$ for $n \in \mathbf{N}$.

(4) We compute that $d_0 = 0$ and $d_1 = 1$. Then for $n \geq 2$, we may instantiate

$$d_n = 1 + 2 \sum_{i=0}^{n-1} (-1)^{n-i} d_i \quad (8)$$

and

$$d_{n-1} = 1 + 2 \sum_{i=0}^{n-2} (-1)^{n-1-i} d_i \quad (9)$$

This time, *adding* the recurrences (8) + (9) turns out to be more helpful as it yields

$$d_n + d_{n-1} = 2 - 2 d_{n-1}$$

and thus

$$d_n = 2 - 3 d_{n-1}.$$

To simplify telescoping, we rearrange this to

$$d_n - \frac{1}{2} = -3 \left(d_{n-1} - \frac{1}{2} \right)$$

and then use the substitution

$$f_n := d_n - \frac{1}{2}$$

from which

$$f_n = -3 f_{n-1}.$$

Telescoping now immediately yields

$$f_n = f_1 (-3)^{n-1},$$

thus

$$f_n = \frac{1}{2} (-3)^{n-1}$$

and so undoing the substitution we end up with

$$d_n = \frac{1}{2} (-3)^{n-1} + \frac{1}{2}.$$

In conclusion, $d_0 = 0$ and $d_n = \frac{1}{2}(1 + (-3)^{n-1})$ for $n \in \mathbf{N}$.

(5) For $n \geq 1$, we have the recurrence

$$e_n = 1 + ne_{n-1}.$$

It is convenient to divide this recurrence by $n!$ as then

$$\frac{e_n}{n!} = \frac{1}{n!} + \frac{e_{n-1}}{(n-1)!}$$

is a simple recurrence for the series $e_n/n!$. Telescoping it yields

$$\frac{e_n}{n!} = \frac{1}{n!} + \frac{1}{(n-1)!} + \dots + \frac{1}{1!} + \frac{e_0}{0!} = \sum_{i=0}^n \frac{1}{i!}$$

for all $n \geq 1$. Therefore,

$$e_n = \left(\sum_{i=0}^n \frac{1}{i!} \right) n!$$

for all $n \in \mathbf{N}_0$ (note that by convention, $0! = 1$).

The above expression may be explicit but it still involves a sum, so we should routinely ask whether there is way to simplify it. The expression in the sum of course makes us think of the function $\exp(\cdot)$. In fact, we know that

$$\sum_{i=0}^{\infty} \frac{1}{i!} = e = 2.71\dots,$$

and thus in the case of our sequence, $e_n < en!$. Due to the nature of the recursion, however, e_n is always an integer, thus we also have $e_n \leq \lfloor en! \rfloor$. Let us compare how close this bound is to the truth.

n	0	1	2	3	4
$e n!$	2.71...	2.71...	5.44...	16.31...	65.24...
$\lfloor en! \rfloor$	2	2	5	16	65
e_n	1	2	5	16	65

So it seems the bound is tight starting $n = 1$. And indeed, if we check,

$$en! - e_n = n! \sum_{i=n+1}^{\infty} \frac{1}{i!} = \frac{1}{n+1} + \frac{1}{(n+1)(n+2)} + \dots$$

is decreasing in n . Since it is below 1 for $n = 1$, it stays below 1 for all n .

We have established

$$e_n = \begin{cases} 1, & \text{if } n = 0; \\ \lfloor en! \rfloor, & \text{otherwise.} \end{cases}$$

Solution 5: Descendants of the Smallest Key

Variante 1: Computation via conditioning on the rank of the root. The usual way, we first obtain

$$\mathbf{E}[S_n] = \sum_{i=1}^n \underbrace{\mathbf{E}[S_n \mid \text{rk}(\text{root}) = i]}_{(*)} \cdot \underbrace{\Pr[\text{rk}(\text{root}) = i]}_{\frac{1}{n}}$$

where

$$(*) = \begin{cases} n, & \text{if } i = 1, \\ \mathbf{E}[S_{i-1}], & \text{otherwise} \end{cases}$$

Denote $s_n := \mathbf{E}[S_n]$. Then this yields a recurrence of the form.

$$s_n = \frac{1}{n} \left(n + \sum_{i=2}^n s_{i-1} \right),$$

holding for all $n \geq 1$. As we are, by now, proficient in solving recurrences, let us multiply by n and then instantiate the recurrence for both n and $n - 1$ so that for $n \geq 1$ we have

$$ns_n = n + \sum_{i=1}^{n-1} s_i \tag{10}$$

and for $n \geq 2$, we get

$$(n-1)s_{n-1} = n-1 + \sum_{i=1}^{n-2} s_i \tag{11}$$

Then subtracting (11) from (10), we obtain

$$ns_n - (n-1)s_{n-1} = 1 + s_{n-1}.$$

Rearranging and dividing by n , this yields

$$s_n = \frac{1}{n} + s_{n-1}.$$

We are familiar with this recursion and know that telescoping it out will produce $s_n = H_n$.

Variante 2: Computation via indicator variables. Alternatively, we can use the well-known indicator variables

$$A_i^j := [\text{node } j \text{ is an ancestor of node } i]$$

In that case we obviously have

$$\begin{aligned} S_n &= \sum_{i=1}^n A_i^1 \\ \Rightarrow s_n = \mathbf{E}[S_n] &= \mathbf{E} \left[\sum_{i=1}^n A_i^1 \right] = \sum_{i=1}^n \mathbf{E}[A_i^1]. \end{aligned}$$

Those expectations have been computed in the lecture notes where we have obtained that

$$\mathbf{E}[A_i^j] = \frac{1}{|i-j|+1}$$

and thus

$$\Rightarrow \mathbf{E}[A_i^1] = \frac{1}{i-1+1} = \frac{1}{i}.$$

Therefore, this variant, too, yields

$$s_n = \sum_{i=1}^n \mathbf{E}[A_i^1] = \sum_{i=1}^n \frac{1}{i} = H_n.$$