

## Matchings

---

A **matching** is a set of (non-loop) edges with no shared endpoints. The vertices incident to an edge of a matching  $M$  are **saturated** by  $M$ , the others are **unsaturated**. A **perfect matching** of  $G$  is matching which saturates all the vertices.

*Examples.*  $K_{n,m}$ ,  $K_n$ , Petersen graph,  $Q_k$ ; graphs without perfect matching

A **maximal matching** cannot be enlarged by adding another edge.

A **maximum matching** of  $G$  is one of maximum **size**.

*Example.* Maximum  $\neq$  Maximal

## Characterization of **maximum** matchings\_\_\_\_\_

Let  $M$  be a matching. A path that alternates between edges in  $M$  and edges not in  $M$  is called an  **$M$ -alternating path**.

An  $M$ -alternating path whose endpoints are unsaturated by  $M$  is called an  **$M$ -augmenting path**.

**Theorem**(Berge, 1957) A matching  $M$  is a maximum matching of graph  $G$  **iff**  $G$  has no  $M$ -augmenting path.

*Proof.* ( $\Rightarrow$ ) Easy.

( $\Leftarrow$ ) Suppose there is no  $M$ -augmenting path and let  $M^*$  be a matching of maximum size.

**What is then  $M \Delta M^*$ ???**

**Lemma** Let  $M_1$  and  $M_2$  be matchings of  $G$ . Then each connected component of  $M_1 \Delta M_2$  is a path or an even cycle.

For two sets  $A$  and  $B$ , the symmetric difference is  $A \Delta B = (A \setminus B) \cup (B \setminus A)$ .

## Hall's Condition and consequences\_\_\_\_\_

**Theorem** (Marriage Theorem; Hall, 1935) Let  $G$  be a bipartite (multi)graph with partite sets  $X$  and  $Y$ . Then there is a matching in  $G$  saturating  $X$  iff  $|N(S)| \geq |S|$  for every  $S \subseteq X$ .

*Proof.* ( $\Rightarrow$ ) Easy.

( $\Leftarrow$ ) Not so easy. Find an  $M$ -augmenting path for any matching  $M$  which does not saturate  $X$ .

(Let  $U$  be the  $M$ -unsaturated vertices in  $X$ . Define

$$T := \{y \in Y : \exists M\text{-alternating } U, y\text{-path}\},$$

$$S := \{x \in X : \exists M\text{-alternating } U, x\text{-path}\}.$$

Unless there is an  $M$ -augmenting path,  $SUU$  violates Hall's condition.)

**Corollary.** (Frobenius (1917)) For  $k > 0$ , every  $k$ -regular bipartite (multi)graph has a perfect matching.

## Graph parameters — Definitions and simple properties

---

The size of the **largest independent set** in  $G$  is denoted by  $\alpha(G)$ .

The size of the **largest matching** (*independent set of edges*) in  $G$  is denoted by  $\alpha'(G)$ .

A **vertex cover** of  $G$  is a set  $Q \subseteq V(G)$  that contains at least one endpoint of every edge. (The vertices in  $Q$  *cover*  $E(G)$ ).

The size of the **smallest vertex cover** in  $G$  is denoted by  $\beta(G)$ .

**Claim.**  $\beta(G) \geq \alpha'(G)$ .

An **edge cover** of  $G$  is a set  $L$  of edges such that every vertex of  $G$  is incident to some edge in  $L$ .

The size of the **smallest edge cover** in  $G$  is denoted by  $\beta'(G)$ .

**Claim.**  $\beta'(G) \geq \alpha(G)$ .

## Certificates

---

Suppose we knew that in some graph  $G$  with 1121 edges on 200 vertices, a particular set of 87 edges is (one of) the largest matching one could find. How could we convince somebody about this?

Once the particular 87 edges are shown, it is easy to check that they are a matching, indeed.

But why isn't there a matching of size 88? Verifying that none of the  $\binom{1121}{88}$  edgesets of size 88 forms a matching could take some time...

If we happen to be so lucky, that we are able to exhibit a vertex cover of size 87, we are saved. It is then reasonable to check that all 1121 edges are covered by the particular set of 87 vertices.

Exhibiting a vertex cover of a certain size **proves** that no larger matching can be found.

## Min-max theorems for bipartite graphs\_\_\_\_\_

**Theorem.** (König (1931), Egerváry (1931)) If  $G$  is bipartite then  $\beta(G) = \alpha'(G)$ .

*Proof.* For any minimum vertex cover  $Q$ , apply Hall's Condition to match  $Q \cap X$  into  $Y \setminus Q$  and  $Q \cap Y$  into  $X \setminus Q$ .

**Lemma.** Let  $G$  be any graph.  $S \subseteq V(G)$  is an independent set iff  $\overline{S}$  is a vertex cover.

Hence  $\alpha(G) + \beta(G) = n(G)$ .

*Proof.* Easy.

**Theorem.** (Gallai, 1959) Let  $G$  be any graph without isolated vertices. Then  $\alpha'(G) + \beta'(G) = n(G)$ .

**Corollary.** (König, 1916) Let  $G$  be a bipartite graph with no isolated vertices. Then  $\alpha(G) = \beta'(G)$ .

*Proof.* Put together the previous three statements.

How to find a **maximum matching** in **bipartite** graphs?\_\_\_\_\_

## **Augmenting Path Algorithm**

**Input.** A **bipartite graph**  $G$  with partite sets  $X$  and  $Y$ ,  
a **matching**  $M$  in  $G$ ,  
the set  $U$  of **unsaturated vertices** in  $X$ .

**Output.** EITHER an  $M$ -augmenting path OR a certificate (a cover of the same size) that  $M$  is maximum.

**Idea.** **Explore**  $M$ -alternating paths from  $U$ , letting  $S \subseteq X$  and  $T \subseteq Y$  be the sets of vertices reached. **Mark** vertices of  $S$  that have been **explored** for path extensions. As a vertex is reached, record the vertex from which it is reached.

**Initialization.**  $S = U$  and  $T = \emptyset$ .

## Iteration.

IF all vertices in  $S$  are **marked** THEN

**stop** and **report** that  $M$  is a **maximum matching** and  $T \cup (X \setminus S)$ , is a **cover of the same size**.

ELSE

select an **unmarked**  $x \in S$  and **explore** its neighbors  $y \in N(x)$ , for which  $xy \notin M$ .

IF  $y$  is unsaturated, THEN

**stop** and **report** an  $M$ -augmenting path from  $U$  to  $y$ .

ELSE

$\exists w \in X$  with  $yw \in M$ . **Update**

$T := T \cup \{y\}$  ( $y$  is reached from  $x$ ),




$S := S \cup \{w\}$  ( $w$  is reached from  $y$ ).

After **exploring** all neighbors of  $x$ , **mark**  $x$  and **iterate**.

**Theorem.** Repeatadly applying the Augmenting Path Algorithm to a bipartite graph produces a maximum matching and a minimum vertex cover.

If  $G$  has  $n$  vertices and  $m$  edges, then this algorithm finds a maximum matching in  $O(nm)$  time.





## Proof of correctness

---

If Augmenting Path Algorithm does what it supposed to, then after at most  $n/2$  application we can produce a maximum matching.

Why does the APA terminate? It touches each edge at most once. Hence running time is  $O(nm)$ .

*What if an  $M$ -augmenting path is returned?* It is OK, since  $y$  is an unsaturated neighbor of  $x \in S$ , and  $x$  can be reached from  $U$  on an  $M$ -alternating path.

*What if the APA returns  $M$  as maximum matching and  $T \cup (X \setminus S)$  as minimum cover?*

Then all edges leaving  $S$  were explored, so there is **no edge between  $S$  and  $Y \setminus T$** .

- Hence  $T \cup (X \setminus S)$  is indeed a cover.
- $|M| = |T| + |X \setminus S|$  (By selection of  $S$  and  $T$ .)

If a cover and a matching have the same size in any graph, then they are both optimal.

$$|M| \leq \alpha'(G) \leq \beta(G) \leq |T \cup (X \setminus S)| = |M|.$$