

Institut für Theoretische Informatik
Dr. Tibor Szabó and Yoshio Okamoto

February 4, 2004

Graph Theory

Problem Set 14

Course Webpage: <http://www.ti.inf.ethz.ch/ew/courses/GT03/>

Due Date: February 11, 2004

Exercise 14.1

(Exercise 6.3.16 in the Textbook)

(-) Prove that the 4-dimensional cube Q_4 is nonplanar. Decompose it into two isomorphic planar graphs.

Exercise 14.2

(Exercise not in the Textbook)

Prove that every outerplanar graph is 3-colorable following each of the ideas below.

- Use Four Color Theorem.
- Use Dirac's theorem.
- Show every simple outerplanar graph is 2-degenerate.

Exercise 14.3

(Exercise not in the Textbook)

In a graph G , to **contract** an edge e with endpoints u, v is to replace u and v with a single vertex whose incident edges are the edges other than e that were incident to u or v . (Hence, the resulting graph has one less edge than G .) A graph H is a **minor** of G if a copy of H can be obtained from G by deleting and/or contracting edges of G , and/or by deleting vertices of G . In such a case we say that G **contains an H -minor**.

Show that the Petersen graph contains a K_5 -minor and a $K_{3,3}$ -minor.

Exercise 14.4

(Exercise 6.2.12 in the Textbook)

(!) Wagner [1937] proved that the following condition is necessary and sufficient for a graph G to be planar: G contains neither K_5 -minor nor $K_{3,3}$ -minor.

- Show that deletion and contraction of edges preserve planarity. Conclude from this that Wagner's condition is necessary.
- Use Kuratowski's Theorem to prove that Wagner's condition is sufficient.

Exercise 14.5

(Exercise 6.3.14 in the Textbook)

(+) Prove that a maximal planar graph is 3-colorable if and only if it is Eulerian. (Hint: For sufficiency, use induction on $n(G)$. Choose an appropriate pair or triple of adjacent vertices to replace with appropriate edges.) (Heawood [1898])

Exercise 14.6

(Exercise not in the Textbook)

A **caterpillar** is a tree in which a single path, called the **spine**, is incident to (or contains) every edge. Prove that in every infinite family of caterpillars there exists two such that one is a minor of the other. (This proves Graph Minor Theorem for caterpillars.)

